Delayed ejaculation



What is delayed ejaculation?

Ejaculation is associated with orgasm but they are two separate physiological processes. While it's possible to reach orgasm and not ejaculate, <u>you can't ejaculate if you don't reach orgasm</u>

Delayed ejaculation is the term commonly used for conditions that result in men experiencing difficulties with reaching orgasm and ejaculating.

We don't really know how many men have delayed ejaculation, but it's probably somewhere between 1 in 5 and 1 in 50.

Delayed ejaculation is more likely to occur if you have particular health conditions or take specific types of medicines.

Symptoms of delayed ejaculation

Delayed ejaculation is diagnosed if you take a long time to reach orgasm or do not reach orgasm at all, even though you want to and are trying, and you are upset or having relationship problems because of it.

Causes of delayed ejaculation

Orgasm and ejaculation are both controlled by the nervous system, so psychological and physical things that affect the necessary nerve activity can both <u>contribute to delayed</u> ejaculation.

Examples of psychological reasons for delayed ejaculation include:

- Performance anxiety
- Religious conflict
- Fears of pregnancy, intimacy, or abandonment.

Physical causes of delayed ejaculation can include:

- Abnormalities in reproductive system development
- Nerve damage from diabetes or surgery
- Aging
- Hormone problems.

The way some men masturbate can affect their ability to <u>reach orgasm and ejaculation</u> with their sexual partners.

Antidepressants and medications to help control blood pressure are among the <u>drugs</u> that may delay ejaculation.

The more medical problems you have, the more likely you are to have delayed ejaculation. Stress and being tired also increase your chance of delayed ejaculation.

Delayed ejaculation



Diagnosis of delayed ejaculation

The reasons for delayed ejaculation vary from person to person and can be quite complicated. Your doctor may need to perform a physical examination, ask some questions and run some tests to work out the best way to treat your delayed ejaculation.

Treatment of delayed ejaculation

There are a variety of medications that can be used to treat delayed ejaculation, but vibratory stimulation of the penis and counselling are usually effective.

If the cause of your delayed ejaculation is psychological, then speaking to someone like a sexual health therapist can help.

Health effects of delayed ejaculation

Delayed ejaculation can make you depressed or anxious and can have negative effects on your relationship with your partner.

Delayed ejaculation can be successfully treated but, because it's such a complex problem, it's difficult to know how long it might take and exactly how successful your treatment will be.

What to do about delayed ejaculation

If you sometimes take a long time to reach orgasm and ejaculate but it doesn't bother you or your partner, then you probably don't need to worry too much about it. But it might still be worth mentioning to your doctor, so they can keep a full record of your health and wellbeing.

If problems with orgasm or ejaculation are bothering you, it's best to see your doctor sooner rather than later.

Questions to ask your doctor about delayed ejaculation

- Could my delayed ejaculation be caused by any medications I'm taking?
- Do you think it's worthwhile for me to talk to a sexual health specialist about my delayed ejaculation?
- Do you have any more questions for me to help you better understand what might be causing my delayed ejaculation?

Medically reviewed by
Dr Michael Lowy
Updated on 23 March 2021